arXiv:1202.3197v1 [cond-mat.quant-gas] 15 Feb 2012

Analytical solutions to the spinor Bose-Einstein condensates

Shi-Jie Yang*,[1] Zhi-Hai Zhang,[1] and Shiping Feng[1]

[1] *Department of Physics, Beijing Normal University, Beijing 100875, China*

We propose a method to analytically solve the one-dimensional coupled nonlinear Gross-Pitaevskii equations which govern the motion of the spinor Bose-Einstein condensates. Classes of exact stationary solutions for the $F = 1$ and $F = 2$ condensates are obtained, from which the soliton complexes are naturally produced and classified. The spin polarizations exhibit parallel or kinked spin density waves. Our method is also applicable to other coupled nonlinear systems in quantum as well as in classical physics.

PACS numbers: 03.75.Mn, 67.85.Fg, 03.75.Lm, 03.75.Hh

Most Bose-Einstein condensates (BECs) of dilute atomic gases realized thus far have internal degrees of freedom arising from spins. The spinor BECs have been realized in the optical traps which confine the atoms regardless of their spin hyperfine states. The direction of the spin can change dynamically due to collisions between the atoms[1–3]. The spinor BECs exhibit a rich variety of magnetic phenomena. They give rise to phenomena that are not present in the single component BECs, including magnetic crystallization, spin textures as well as fractional vortices[4–6]. Moreover, there exists an interplay between superfluidity and magnetism due to the spin-gauge symmetry. A ferromagnetic BEC spontaneously creates a supercurrent as the spin is locally rotated[7, 8].

A spinor condensate formed by atoms with spin $F = 1$ is described by a macroscopic wave function with three components $\Psi = (\psi_{+1}, \psi_0, \psi_{-1})^T$. The mean-field Hamiltonian is expressed as[9, 10]

$$H = \int d\mathbf{r} \quad \{\sum_{m=-1}^{1} \psi_m^* [-\frac{\hbar^2}{2M} \bigtriangledown^2 + V(\mathbf{r})]\psi_m + \frac{\bar{c}_0}{2} n_{tot}^2 + \frac{\bar{c}_2}{2} |\mathbf{F}|^2\}, \tag{1}$$

where the spin-polarization vector $\mathbf{F} = \psi_m^* \hat{\mathbf{F}}_{mn} \psi_n$ with $\hat{\mathbf{F}}^i$ $(i = x, y, z)$ the rotational matrices. The coupling constants $\bar{c}_0 = (g_0 + 2g_2)/3$ and $\bar{c}_2 = (g_2 - g_0)/3$, with g_F relating to the s-wave scattering length of the total spin-F channel as $g_F = 4\pi\hbar^2 a_F/M$. $n_{tot} = |\psi_1|^2 + |\psi_0|^2 + |\psi_{-1}|^2$ and $V(\mathbf{r})$ is the external potential. The full symmetry of the Hamiltonian is $SO(3) \times U(1)$. In the ground state, the symmetry is spontaneously broken in several different ways, leading to a number of possible phases[9–12]. The energies are degenerate for an arbitrary state Ψ and its globally spin-rotational states $\tilde{\Psi} = U\Psi$, where U is the three-dimensional rotational matrix in the spin space which is expressed by the Euler angles as $U(\alpha, \beta, \gamma) = e^{-i\hat{F}_z\alpha} e^{-i\hat{F}_y\beta} e^{-i\hat{F}_z\gamma}$.

We are concerned with the quasi-one-dimensional (1D) $F = 1$ and $F = 2$ BEC in a uniform external potential ($V(\mathbf{r}) = 0$). The dynamical motions of the $F = 1$ spinor wavefunctions are governed by $i\partial_t\psi_m = \delta H/\delta\psi_m^*$, which are explicitly written as the coupled nonlinear Gross-Pitaevskii equations (GPEs),

$$i\partial_t\psi_m = [-\frac{\hbar^2}{2M}\partial_x^2 + c_0 n_{tot} + c_2(\Psi^\dagger \hat{F}^i \Psi)\hat{F}_{mn}^i]\psi_n, \tag{2}$$

where $c_0 = \bar{c}_0/2a_\perp^2$ and $c_2 = \bar{c}_2/2a_\perp^2$ are the reduced coupling constants with $a_\perp$ the transverse width of the quasi-1D system. There is a lot of theoretical works that numerically solve Eqs.(2) or the corresponding stationary equations[13–15]. Nevertheless, it is of great interest if analytical solutions can be acquired since they provide us with further insights into the correlations of the condensates and help to explore the possible topological structures in details. There are some works that utilized the similar transformation to obtain the soliton solutions for systems with the assumption of time or spatial variations in the coupling constant[16–19]. Various approximations are employed to study the solitons such as bright and dark solitons in the $F = 1$ spinor BECs[20–23]. However, the general exact solutions for the spinor BECs, especially for the $F = 2$ system, are still absent in literature. Attempts in this direction are usually frustrated by the tremendous complexity of the coupled nonlinear GPEs. The challenges are two-fold: one is the nonlinear density-density interactions which is associated with the c_0 term, while the other is the spin-exchange couplings between the hyperfine states which is associated with the c_2 term.

In this Letter we propose a general method which simultaneously decouples the nonlinear density-density interactions and the spin-spin interactions in the GPEs (2). We obtain classes of the exact stationary solutions without any approximations or the assumptions of time or spatial variations of the coupling strengths. The solutions are either in real or in complex forms and are expressed by combinations of the Jacobi elliptical functions. We found that the spin-polarizations exhibit either parallel or kinked spin density-wave configurations. The polar states ($|\mathbf{F}| = 0$) with non-uniform atomic density are

---

*Corresponding author: yangshijie@tsinghua.org.cn

also obtained. The stationary soliton or the soliton train complexes are naturally produced, which are identified as polarized or vector solitons, and polar or scalar solitons, respectively. Furthermore, we obtain new and more general stationary solutions by rotating the states in the spin space. All solutions have been numerically verified to be stationary by the dynamical evolution with Eqs.(2).

First we deal with the $F = 1$ System. For convenience we choose in the following discussions $\hbar = M = 1$ as the units. c_0 and c_2, or a_0 and a_2, are treated as free parameters. The stationary GPEs (2) are written as

$$\begin{aligned}
\mu\psi_1 &= [-\frac{1}{2}\partial_x^2 + (c_0+c_2)(|\psi_1|^2+|\psi_0|^2) + (c_0-c_2)|\psi_{-1}|^2]\psi_1 + c_2\psi_0^2\psi_{-1}^*, \\
\mu\psi_0 &= [-\frac{1}{2}\partial_x^2 + (c_0+c_2)(|\psi_1|^2+|\psi_{-1}|^2) + c_0|\psi_0|^2]\psi_0 + 2c_2\psi_0^*\psi_1\psi_{-1}, \\
\mu\psi_{-1} &= [-\frac{1}{2}\partial_x^2 + (c_0+c_2)(|\psi_{-1}|^2+|\psi_0|^2) + (c_0-c_2)|\psi_1|^2]\psi_{-1} + c_2\psi_0^2\psi_1^*.
\end{aligned} \tag{3}$$

The periodic boundary conditions,

$$\psi_m(1) = \psi_m(0), \qquad \psi_m'(1) = \psi_m'(0). \tag{4}$$

is adopted in our deduction. In order to seek the analytical solutions we decouple spin-spin interactions in the Eqs.(3) by assuming $\psi_{-m}^* = \delta\psi_m$, with $\delta = \pm 1$. It is easy to show that $\delta = +1$ corresponds to (partially) spin-polarized states ($|\mathbf{F}| \neq 0$) whereas $\delta = -1$ corresponds to spin-unpolarized or polar states ($|\mathbf{F}| = 0$). At the same time, the nonlinear density-density interactions between different components are decoupled by using the unique properties of the Jacobi elliptical functions. There are real and complex forms of solutions which are explicitly described as follows.

Real solutions. Consider real solutions of the *sn*-*cn*-*sn* form

$$\begin{pmatrix} \psi_1(x) \\ \psi_0(x) \\ \psi_{-1}(x) \end{pmatrix} = \begin{pmatrix} Asn(kx,m) \\ Dcn(kx,m) \\ \delta Asn(kx,m) \end{pmatrix}, \tag{5}$$

where sn, cn, (or dn below) are the Jacobi elliptical functions and $k = 4jK(m)$ with m the modulus $(0 < m < 1)$. A and D are the real amplitudes. In the context we always take the number of periods $j = 2$ as examples. One has

$$|\psi_{-1}|^2 = |\psi_1|^2, \quad |\psi_0|^2 = D^2 - \frac{D^2}{A^2}|\psi_1|^2. \tag{6}$$

Substituting (6) into the stationary equations (3), we obtain three decoupled equations

$$\tilde{\mu}_m\psi_m = -\frac{1}{2}\psi_m'' + \tilde{\gamma}_m|\psi_m|^2\psi_m, \quad (m = 0, \pm 1) \tag{7}$$



FIG. 1: (color) (a) The density distribution of the stationary solutions (5) for the $F = 1$ condensates with $\delta = +1$. $m = 0.5$ and $\mu = 600$. (c) The spin-rotational solution of (5). (b, d) are respectively the corresponding soliton solutions with $m \to 1$ and $\mu = 8000$.

where the effective chemical potentials $\tilde{\mu}_m$ and interacting constants $\tilde{\gamma}_m$ are defined as

$$\begin{aligned}
\tilde{\mu}_{\pm 1} &= \mu - (c_0 + c_2 + c_2\delta)D^2 \\
\tilde{\mu}_0 &= \mu - 2(c_0 + c_2 + c_2\delta)A^2 \\
\tilde{\gamma}_{\pm 1} &= 2c_0 - (c_0 + c_2 + c_2\delta)\tfrac{D^2}{A^2} \\
\tilde{\gamma}_0 &= c_0 - 2(c_0 + c_2 + c_2\delta)\tfrac{A^2}{D^2}.
\end{aligned} \tag{8}$$

The effective chemical potentials $\tilde{\mu}_m$ and amplitudes



FIG. 2: (color) (a) The density distribution of the stationary solutions of *cn*-*sn*-*cn* form for the $F = 1$ condensates with $\delta = -1$. $m = 0.5$, $\mu = 6000$. (c) is the spin-rotational solution. (b,d) are respectively the corresponding soliton solutions with $m \to 1$, $\mu = 80000$.



FIG. 3: (color online) The phase distribution for the $F = 1$ complex solutions. (a) $n = 1$ with $m = 0.29$. (b) $n = 2$ with $m = 0.92$.

A, D can be obtained self-consistently as,

$$\begin{aligned} &\tilde{\mu}_{\pm 1} = \tfrac{1}{2}k^2(1+m^2) \\ &\tilde{\mu}_0 = \tfrac{1}{2}k^2(1-2m^2) \\ &A^2 = \tfrac{m^2k^2}{\tilde{\gamma}_1} = \tfrac{\mu - \frac{1}{2}k^2(1-2m^2)}{2(c_0+c_2)+2c_2\delta} \\ &D^2 = -\tfrac{m^2k^2}{\tilde{\gamma}_0} = \tfrac{\mu-\frac{1}{2}k^2(1+m^2)}{(c_0+c_2)+c_2\delta}. \end{aligned} \tag{9}$$

Formula (9) implies that the effective nonlinear interactions for $\psi_\pm$ must be repulsive ($\tilde{\gamma}_{\pm 1} > 0$) while for ψ_0 attractive ($\tilde{\gamma}_0 < 0$) which imposes constraints on the coupling strengths c_0 and c_2. Here the value of m is arbitrary.

More general solutions are obtained from solution (5) by making use of the $SO(3)$ rotational symmetry in the spin space $\tilde{\psi}_m = U_{mn}\psi_n$. It is clear that $\tilde{\psi}_m$ are also stationary solution. In the following we fix the three Euler angles $\alpha = 7\pi/13, \beta = \pi/4, \gamma = \pi/11$ as examples.

Figure 1 display the density distribution of the *sn*-*cn*-*sn* solution with $\delta = +1$, which is a (partially) spin-polarized stationary states. The values of parameters a_0 and a_2 are indicated in the figures. Fig.1(a) illustrates the solution (5) and Fig.1(c) is the result after rotation of the solution (5) in the spin space, where $m = 0.5$ and $\mu = 600$. We note that ψ_{+1} and ψ_{-1} are staggered after the spin-rotation. The soliton solutions are naturally produced by taking the limit $m \to 1$. Fig.1(b) exhibits a typical dark-bright-dark soliton ($\mu = 8000$). The spin-rotating solution in Fig.1(d) reveals more complicated soliton structure. Single soliton solutions can be obtained by directly substituting the Jacobi elliptical functions in Eq.(5) with the hyperbolic functions. Other soliton solutions such as bright-dark-bright complexes can be constructed by the same way. These solitons, being (partially) spin-polarized ($|\mathbf{F}| \neq 0$), may be properly called polarized or vector solitons.

Figure 2 display the stationary solution of the *cn*-*sn*-*cn* form with $\delta = -1$, which is a polar state with spin-polarization $|\mathbf{F}| = 0$. It should be noted that ψ_{+1} and ψ_{-1} remain overlapped after the spin-rotation. In this case, the soliton limit exhibits bright-dark-bright complexes. In comparison to the polarized solitons, these spin-unpolarized solitons are called polar or scalar solitons.

Complex solutions. We assume $\psi^*_{-1} = \psi_1$ ($\delta = +1$) and consider the following *cn*-*sn*-*cn* form of complex solution,

$$\begin{pmatrix} \psi_1 \\ \psi_0 \\ \psi_{-1} \end{pmatrix} = \begin{pmatrix} f(x)e^{i\theta(x)} \\ Dsn(kx,m) \\ f(x)e^{-i\theta(x)} \end{pmatrix}, \tag{10}$$

where $f(x) = \sqrt{A + Bcn^2(kx,m)}$. A, B and D are real constants. One has

$$|\psi_{-1}|^2 = |\psi_1|^2, \quad |\psi_0|^2 = D^2(1+\frac{A}{B}) - \frac{D^2}{B}|\psi_1|^2. \tag{11}$$

Substituting (11) into the coupled GPEs (3), we again obtain the decoupled equations (7) but with a different set of effective chemical potentials and interaction con-



FIG. 4: (color) The density distribution of each hyperfine state for the $F = 1$ complex solutions. Left column are solutions from Eq.(10) with (a) $n = 1, m = 0.29, \mu = 90$, (c) $n = 2, m = 0.92, \mu = 800$, and (e) $n = 3, m = 0.99, \mu = 2000$, respectively. Right column are the corresponding rotating solutions.



FIG. 5: (color online) The spin-polarization distribution for the $F = 1$ stationary solutions. (a) is the real solution of Eq.(5) with $\delta = +1$. (b)-(d) are the complex solutions of Eq.(10) for $n = 1$, $n = 2$, and $n = 3$, respectively.

stants,

$$\begin{array}{l}\tilde{\mu}_{\pm1} = \mu - (c_0 + 2c_2)\frac{D^2}{B}(A + B)\\ \tilde{\mu}_0 = \mu - 2(c_0 + 2c_2)(A + B)\\ \tilde{\gamma}_{\pm1} = 2c_0 - (c_0 + 2c_2)\frac{D^2}{B}\\ \tilde{\gamma}_0 = c_0 - 2(c_0 + 2c_2)\frac{B}{D^2}\end{array} \qquad (12)$$

The complex form of solutions to the decoupled Eqs.(7) are self-consistently solved as

$$\begin{array}{l}B = -\frac{m^2k^2}{\tilde{\gamma}_1}\\ A = \frac{1}{3\tilde{\gamma}_1}(2\tilde{\mu}_1 - (1 - 2m^2)k^2)\\ D^2 = \frac{m^2k^2}{\tilde{\gamma}_0}\end{array} , \qquad (13)$$

which leads to

$$\begin{array}{l}D^2 = -2B = \frac{m^2k^2}{2(c_0+c_2)}\\ \tilde{\gamma}_1 = 2\tilde{\gamma}_0 = 4(c_0 + c_2).\end{array} \qquad (14)$$

One observes that the effective interactions $\tilde{\gamma}_m > 0$ for this form of complex solution. The phase is

$$\theta(x) = \int_0^x \frac{\alpha_1}{f^2(\xi)} d\xi, \qquad (15)$$

where $\alpha_1 = \pm(2\tilde{\mu}_1 A^2 - 2\tilde{\gamma}_1 A^3 + k^2(1 - m^2)AB)^{\frac{1}{2}}$ is an integral constant. The periodic boundary conditions (4) require that the amplitudes and phase satisfy

$$f(1) = f(0), \quad \theta(1) - \theta(0) = 2j\pi \times n, \qquad (16)$$

respectively, where n is an integer. The periodic condition for the phase is satisfied by adjusting the modulus m of the Jacobi elliptical functions.

Figure 3 display the phase distribution for (a) $n = 1$ and (b) $n = 2$, respectively. They exhibit a four-step staircase structure for $j = 2$. Figure 4 show the density distribution for each hyperfine state of the complex stationary solution (10) and its rotational state for (a, b) $n = 1$, ($m = 0.29, \mu = 90$) (c, d) $n = 2$ ($m = 0.92, \mu = 800$) and (e, f) $n = 3$ ($m = 0.99, \mu = 2000$), respectively. In contrast to the real solutions, the complex solutions bear mass counter-flows as well as spin currents, yet the density distributions remain stationary.

Figure 5 show the spin-polarization vector distribution $\mathbf{F}(x)$ of the above stationary solutions. Fig.5(a) is for the real solution (5) which exhibits a parallel polarization configuration similar to a spin-density-wave. Fig.5(b-d) are for the complex solutions (10) with $n = 1$, $n = 2$ and $n = 3$, respectively. The spin-polarizations exhibit kinked structures.

This method also is applicable to the $F = 2$ condensates. Figure 6 show the *sn-cn-sn-cn-sn* form of real stationary solutions which correspond to the spin-unpolarized states. The details of the results will be present elsewhere.

In summary, we have developed a method to systematically solve the one-dimensional coupled nonlinear GPEs which govern the motion of the spinor BECs. Several classes of analytical stationary solutions for the $F = 1$ and $F = 2$ condensates are obtained. Obviously, we can also obtain a lot of other stationary solutions with differ-



FIG. 6: (color) The real solutions of the *sn*-*cn*-*sn*-*cn*-*sn* form for the $F = 2$ condensates. (a) $m = 0.5, \mu = 90$. (b) $m \to 1, \mu = 6000$. (c,d) are the corresponding rotating solutions.

ent combinations of the Jacobi elliptical functions, which will be published in another paper. We emphasize the our method is general and exact, without any approximations, and can be extended to other nonlinear systems such as the coupled nonlinear Klein-Gordon equations, the dynamical coupled nonlinear Schrodinger equations, the coupled KdV equations, or the circularly symmetric coupled sexticoscillators, et al. Works in this direction are in progress.

This work is supported by the funds from the Ministry of Science and Technology of China under Grant No. 2012CB821403 and 2011CB921700, and by the National Natural Science Foundation of China under grant No. 10874018.

[1] M. D. Barrett, et al., Phys. Rev. Lett. **87**, 010404 (2001).
[2] A. Gorlitz, et al., Phys. Rev. Lett. **90**, 090401 (2003).
[3] A. E. Leanhardt, et al., Phys. Rev. Lett. **90**, 140403 (2003).
[4] Stenger, J. et al., Nature 396, 345 (1998).
[5] H. J. Miesner, et al., Phys. Rev. Lett. **82**, 2228 (1999).
[6] M. Kobayashi, et al., Phys. Rev. Lett. **103**, 115301(2009).
[7] M. Ueda and Y. Kawaguchia, arXiv: 1001.2072 (unpublished).
[8] Y. Kawaguchi, et al., Phys. Rev. Lett. **96**, 080405 (2006).
[9] T.-L. Ho, Phys. Rev. Lett. **81**, 742 (1998).
[10] M.-S. Chang, et al., Phys. Rev. Lett. **92**, 140403 (2004).
[11] K. Murata, et al., Phys. Rev. A **75**, 013607 (2007).
[12] A. Imambekov, et al., Phys. Rev. A **68**, 063602 (2003).
[13] L. Li, et al., Phys. Rev. A **72**,033611 (2005).
[14] W. Zhang, et al., Phys. Rev. A **75**, 043601 (2007).
[15] H. E. Nistazakis, et al., Phys. Rev. A **77**,033612(2008).
[16] J. Belmonte-Beitia, et al., Phys. Rev. Lett. **98**, 064102 (2007).
[17] G. Theocharis, et al., Phys. Rev. A **72**, 033614 (2005).
[18] A.T.Avelar, et al., Phys. Rev. E **79**, 025602(R) (2009).
[19] D.-S. Wang, et al., Phys. Rev. A **82**, 023612 (2010).
[20] L. D. Carr, et al., Phys. Rev. A **62**, 063610 (2000).
[21] R. M. Bradley, et al., J. Phys. A: Math. Gen. **38**, 1901 (2005).
[22] L. Li, et al., Phys. Rev. E **73**, 066610 (2006).
[23] D. Yan, et al., Phys. Rew. A **84**, 053630 (2011).